



ESB PLUS

WATER SOLUBLE POWDER 100 GRAM

TREATMENT OF COCCIDIOSIS

Coccidiosis is an infectious disease caused by protozoa (small single-celled parasites). It develops in the host's *intestinal cells* and damages the intestinal wall. Digestion is then severely impaired and birds lose weight. Young birds are very susceptible!

Atoxoplasmosis is also a specific form of coccidiosis. This parasite enters the bloodstream during its cycle and can settle in organs such as the liver, spleen, lungs and brain. Birds develop thickened livers, bloody spots on the intestines, wheezing and sometimes loss of balance.

ESB PLUS intervenes in the coccidiosis cycle and prevents further development. It is therefore a coccidiostat; it stops the cycle. It has no negative effects on fertilisation, laying, growth or hatching outcomes.



INDICATIONS

- Preventive treatment before the start of breeding.
- Treatment of coccidiosis infection.
- Control of atoxoplasmosis (Lankesterella).
- Preventive protection during the moulting period.

DOSAGE AND ADMINISTRATION

Oral administration via drinking water

• In case of coccidiosis: 2 g per litre of drinking water for 2 x 5 days with 2 days of rest in between. This treatment can be repeated in the same way after 2 weeks.

 In case of atoxoplasmosis: 1 g per litre of drinking water for 4 periods of 5 days with 2 days rest in between each time.

Change the medicinal water daily and provide it as the only source of drinking water during treatment. Additional vitamins can be given during rest periods.

Each pack contains a measuring spoon of 8 grams.



Ordering through:

WWW.PANTEX-COUTTEEL.COM